

LETTER TO EDITOR

Year : 2014 | Volume : 47 | Issue : 1 | Page : 147–149

Free saphenous venous flap and its modifications: Turbo- and supercharged**Kathleen Yee Kuin Chang, Wan Azman Wan Sulaiman, Arman Zaharil Mat Saad**

Department of Reconstructive Science, School of Medical Sciences, Universiti Sains Malaysia, Kelantan, Malaysia

Correspondence Address:

Wan Azman Wan Sulaiman

Department of Reconstructive Science, School of Medical Sciences, Health Campus, Universiti Sains Malaysia, 16150 Kubang Kerian, Kelantan Malaysia

How to cite this article:

Chang KY, Sulaiman WW, Mat Saad AZ. Free saphenous venous flap and its modifications: Turbo- and supercharged. Indian J Plast Surg 2014;47:147-149

How to cite this URL:

Chang KY, Sulaiman WW, Mat Saad AZ. Free saphenous venous flap and its modifications: Turbo- and supercharged. Indian J Plast Surg [serial online] 2014 [cited 2018 Jul 15];47:147-149

Available from: <http://www.ijps.org/text.asp?2014/47/1/147/129658>**Full Text**

Sir,

Nakayama et al. first reported an experimental work on venous flap in 1981, [1] followed by reports of clinical application and development of the arterialized venous flaps by various authors. [2] The venous flap is useful where arterial donor site morbidity is undesirable, where large healthy veins are found proximal and distal to the wound, or soft-tissue coverage is needed in arterial reconstruction. The disadvantages include higher flap loss rate and persistent oedema. [3] We demonstrated the use of a modification of saphenous venous flow-through flap in the reconstruction of upper limb arteries with soft-tissue coverage. To the best of our knowledge, this is the first reported case using a turbo- and supercharged saphenous venous flap.

Patient B is a 64-year-old gentleman who sustained an open posterior dislocation of right elbow and right brachial artery injury. An emergency right brachial artery repair and external fixation were performed but he subsequently developed a bleeding right brachial artery aneurysm. The patient was planned for resection of aneurysm and reconstruction of right brachial artery with long saphenous vein graft.

Intra-operative findings revealed fragile and inflamed overlying skin with macerated distal portion of brachialis up to point of insertion and macerated proximal portion of the flexor origin. Resection of the pseudoaneurysm measuring 6 cm × 7 cm and debridement of the macerated tissues left a large defect at the medial aspect of the arm and forearm, exposing some tendons and ulnar nerve.

In view of the need for soft-tissue reconstruction besides the brachial artery reconstruction, we initially harvested a left saphenous venous flap with a skin paddle measuring 13.5 cm × 7 cm. Two arterial perforators (proximal and distal) were identified using pencil Doppler prior to flap elevation and were included in the flap design to increase the survival of the skin paddle. These perforators were traced to the left posterior tibial artery [Figure 1]. We eventually harvested a free saphenous venous flow-through flap turbocharged with the posterior tibial artery perforator and supercharged with cephalic vein to reconstruct the brachial artery and surrounding soft-tissue [Figure 2]. Donor site was closed primarily. {Figure 1}{Figure 2}

After establishing flow of the brachial artery using the saphenous vein, we noted strong arterial blood flow through the vena comitans of the proximal perforator (via communication between the deep and superficial venous system of the lower limb). Therefore, the proximal perforator was anastomosed to the vena comitans of the perforator ("arterial limb" of the flap) as

turbocharge [Figure 3]. Post-anastomosis, a strong arterial signal was detected using pencil Doppler. A cutaneous vein from the flap was anastomosed to the cephalic vein (venous limb of the flap) to supercharge the venous outflow of the flap [Figure 4].

The flap survived with the brachial artery and the anastomosis between the vena comitans of the perforator artery and the perforator artery ("arterial limb" of the flap) remaining patent with strong arterial signal when monitored with pencil Doppler for up to a week. Despite the large skin paddle, the recipient site required split-thickness skin graft for complete wound closure [Figure 5].
{Figure 3}{Figure 4}{Figure 5}

A study by Koshima et al. [4] demonstrated that a supercharged arterialized saphenous venous flap measuring 11 cm × 7 cm or more can survive without superficial necrosis. We developed a modification to our initial saphenous venous flow-through flap. Initially the posterior tibial artery perforator was included in the flap with the intention to supercharge the arterial supply of the flap. However in view of the good arterial flow from the vena comitans of the posterior tibial artery after reconstruction of the brachial artery, we decided to turbocharge the flap. The vena comitans received arterial blood in retrograde fashion from the saphenous vein graft via communications between the deep and superficial venous system. The arterial inflow to the skin paddle was established by anastomosing the vena comitans of the perforator to the perforator. We believed that by creating an arterial inflow to the skin paddle of the flap, a more physiologic blood flow was provided. The venous outflow of the skin paddle was supercharged by anastomosing the cutaneous vein of the skin paddle to the cephalic vein. The flap did not experience any venous congestion for the first 72 hours post-operation.

We believed by turbocharging and supercharging the flap, the skin paddle received a more reliable and physiological inflow and outflow, thus increasing its survival. As this modification is a new technique, we have yet to establish the reproducibility of this technique in improving flap survival. However, we have shown that the skin paddle in patient B's flap survived 100%.

The saphenous venous flow-through flap is a functional alternative with minimal donor site morbidity for reconstruction of major limb vessels. The flap can be turbocharged if a perforator artery is discovered supplying the skin paddle and supercharged by creating another venous outflow to improve flap survival.

References

- 1 Nakayama Y, Soeda S, Kasai Y. Flaps nourished by arterial inflow through the venous system: An experimental investigation. *Plast Reconstr Surg* 1981;67:328-34.
- 2 Yoshimura M, Shimada T, Imura S, Shimamura K, Yamauchi S. The venous skin graft method for repairing skin defects of the fingers. *Plast Reconstr Surg* 1987;79:243-50.
- 3 Deune EG, Rodriguez E, Hafez D, Frassica F. Arterialized venous flow-through flap for simultaneous reconstruction of a radial artery defect and palmar forearm soft-tissue loss from sarcoma resection. *J Reconstr Microsurg* 2005;21:85-91.
- 4 Koshima I, Soeda S, Nakayama Y, Fukuda H, Tanaka J. An arterialised venous flap using the long saphenous vein. *Br J Plast Surg* 1991;44:23-6.

Sunday, July 15, 2018

[Site Map](#) | [Home](#) | [Contact Us](#) | [Feedback](#) | [Copyright and Disclaimer](#)